

POPULATIONS OF ATLANTIC RIDLEY SEA TURTLES (LEPIDOCHELYS KEMPI) IN APALACHEE BAY, FLORIDA, COASTAL WATERS

Ann Rudloe¹

Jack Rudloe¹

Larry Ogren²

¹Gulf Specimen Marine Laboratories, Inc., P.O. Box 237, Panacea, Florida 32346 USA

²NOAA/NMFS Panama City Laboratory, 3500 Delwood Beach Road, Panama City, Florida 32408 USA

A total of 106 Kemp's ridley sea turtles, (Lepidochelys kempi), was captured, measured, tagged, and released from 1984-1988 in Wakulla and Franklin counties, Florida. Turtles ranged from 20.3 to 57.9 cm straight line carapace length (mean 36.4 cm), and weights ranged from 635 grams to 13.7 kg. All turtles were immature and were captured in all months of the year. Turtles taken in winter were significantly larger than those taken in summer. Carapace epibionts and mud stains suggested that two individuals had over-wintered. Turtles were taken over sand, mud, and seagrass bottoms and at depths ranging from 30 cm to 32 m. A significant length-depth relationship was observed. The smallest turtles were taken from depths < 9 m. Four recoveries were made, three of which returned to their point of capture upon release. Offshore occurrences during winter months to depths > 30 m for three individuals were observed.



PROCEEDINGS OF THE NINTH ANNUAL WORKSHOP ON SEA TURTLE CONSERVATION AND BIOLOGY

**7-11 February 1989
Jekyll Island, Georgia**

**Compilers:
Scott A. Eckert
Karen L. Eckert
Thelma H. Richardson**

August 1989

The Technical Memorandum Series is used for documentation and timely communication of preliminary results, interim reports, or special-purpose information. The contents have not received complete formal review or editorial control.

**U.S. DEPARTMENT OF COMMERCE
Robert A. Mosbacher, Secretary**

**NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
John A. Knauss, Administrator**

**NATIONAL MARINE FISHERIES SERVICE
James W. Brennan, Assistant Administrator for Fisheries**

Cover Artwork: ©Lloyd Logan 1989